




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, A01H 5/00 // C07K 14/08		A1	(11) International Publication Number: WO 98/07875
(21) International Application Number: PCT BE97/00092		(43) International Publication Date: 26 February 1998 (26.02.98)	
(22) International Filing Date: 18 August 1997 (18.08.97)		(74) Agents: VAN MALDEREN, Eric et al., Office Van Malderen, Place Reine Fabiola 6-1, B-1083 Brussels (BE).	
(30) Priority Data: 96870106.0 19 August 1996 (19.08.96) EP (34) Countries for which the regional or international application was filed: BE et al.		(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, DE (Utility model), EE, GE, HU, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for all designated States except US): SES EUROPE N.V./S.A. [BE/BE], Industriepark 15, B-3300 Tienen (BE).		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(72) Inventors; and (75) Inventors/Applicants (for US only): GUILLEY, Hubert [FR/FR]; 32, rue de l'Herbe, F-67370 Berstett (FR); JONAR, Gérard [FR/FR]; 9, quai de Chanoine Winterer, F-67000 Strasbourg (FR); RICHARDS, Ken [FR/FR]; 2, rue Principale, F-67370 Pfulgriesheim (FR); BOUZOUBAA, Salah [FR/FR]; 11, rue de Berstett, F-67200 Strasbourg (FR); BLEYKASTEN-GROSSHANS, Claudine [FR/FR]; 9, rue du Renard Préchant, F-67000 Strasbourg (FR); WEYENS, Guy [BE/BE]; Dwarsbos 25, B-1650 Beersel (BE); LEFEBVRE, Marc [BE/BE]; Rue de la Chapelle Stevenaert, 75, B-1370 Jodoigne (BE).			

(54) Title: METHOD FOR INDUCING VIRAL RESISTANCE INTO A PLANT

(57) Abstract

The present invention concerns a method for inducing resistance to a virus comprising a TGB3 sequence with the proviso that it is not the potato virus X, into a cell plant or a plant, comprising the following steps: preparing a nucleic acid construct comprising a nucleic acid sequence corresponding to at least 70 % of the nucleic acid sequence of TGB3 of said virus or its corresponding cDNA, being operably linked to one or more regulatory sequence(s) active in a plant, transforming a plant cell with the nucleic acid construct, and possibly regenerating a transgenic plant from the transformed plant cell. The present invention is also related to the plant obtained.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5

METHOD FOR INDUCING VIRAL RESISTANCE INTO A PLANT

10

Field of the invention

The present invention is related to a method for inducing viral resistance into a cell and plant, especially BNYVV-resistance into a sugar beet cell and
15 plant and the viral resistant cell and plant obtained.

Background of the invention and state of the art

The widespread viral disease of the sugar beet plant (Beta vulgaris) called Rhizomania is caused by a
20 furovirus, the beet necrotic yellow vein virus (BNYVV) (23, 24) which is transmitted to the root of the beet by the soilborne fungus Polymyxa betae (25).

The disease affects significantly acreages of the area where the sugar beet plant is grown for industrial
25 use in Europe, USA and Japan and is still in extension in several places in Western Europe (26, 27). As there exists no practical method to effectively control the spread of the virus at a large scale by chemical or physical means (28), neither in the plants nor in the soil, the main focus
30 has been to identify natural sources of resistance within the sugar beet germplasm and to develop by breeding, varieties of sugar plants beet expressing the resistance

genes. A variety of such tolerance genes to the virus has been identified and, some have been successfully used in the breeding of commercial sugar beet varieties (29, 30, 31).

5 Only the use of BNYVV-resistant or tolerant varieties will enable farmers to grow sugar beet plants in BNYVV-infected areas where sugar beet plant is an essential component of the crop rotation and contributes significantly to the grower's income.

10 A number of detailed studies have shown that a difference in susceptibility to the BNYVV-infection among sugar beet genotypes or varieties, generally reflect difference in the diffusion or translocation of the virus in the root tissues (32).

15 However, there are still few reports which indicate clearly that the tolerance genes, even from differing sources of sugar beet germplasm or wild relatives germplasm (33), would provide distinct mechanisms of resistance. Such a situation would represent a more
20 manageable situation to design long lasting BNYVV-resistance strategies.

 Since 1986, number of reports and publications have described the use of isolated viral gene sequences expressed in plant to confer a high level of
25 tolerance against the virus or even to confer a broad spectrum type of resistance against a number of related viruses (34, 35, 36). One of the most documented viral resistance strategy based on genetic engineering, in many cultivated species such as potato, squash, cucumber or
30 tomato, is the use of the viral gene sequence encoding the coat-protein of the target virus (37) which under the control of plant regulatory elements, will be expressed in

the plant.

However in the case of the coat-protein mediated resistance, the expression of a certain level of resistance in the transgenic plant might be attributed to
5 different mechanisms such as RNA co-suppression and not necessarily to the production of the protein sequence.

In general, the virus sequence will be transformed in an appropriated cell or a tissue culture of the plant species using an Agrobacterium mediated
10 transformation system or a direct gene transfer method according to the constraints of the tissue culture or cell culture method which can be successfully applied in a given species. A whole plant will be regenerated and the expression of the transgene will be characterized.

15 Though sugar beet is known as a recalcitrant species in cell culture, limiting the extent of practical genetic engineering applications in that species, there are number of isolated reports of successful transformation and regeneration of whole plants (38). A few examples of
20 engineering tolerance to the BNYVV by transforming and expressing the BNYVV coat-protein sequence in the sugar beet genome have also been published (39, WO91/13159) though they rarely report data on whole functional transgenic sugar beet plants (40). In particular, reports
25 show limited data on the level of resistance observed in infected conditions with transgenic sugar beet plants transformed with a gene encoding a BNYVV coat-protein sequence (41, 42).

A complete technology package including a
30 sugar beet transformation method and the use of the expression of the BNYVV coat-protein sequence as resistance source in the transgenic sugar beet plant obtained by said

transformation method has been described in the Patent Application WO91/13159.

Based on the information published, it can not be concluded that the coat-protein mediated resistance mechanism provides any potential for conferring to the sugar beet plant a total immunity to the BNYVV-infection by inhibiting completely the virus multiplication and diffusion mechanisms. To identify a resistance mechanism which enables to block significantly the spread of the virus at the early stage of the infection process would be a major criteria of success to develop such a transgenic resistance, in addition to the fact that even a level of resistance comparable to those known from the genes of resistance identified within the sugar beet germplasm would diversify the mechanisms of resistance available.

Because the disease is shown to expand in many countries or areas, at a speed depending upon the combination of numerous local environmental and agricultural factors, there is a major interest to diversify the sources of genetic resistance mechanisms which may, alone or in combination, confer a stable and long lasting resistance strategy in the current and future varieties of sugar beet plants which are grown for industrial use.

The publication of Xu H. et al. (Plant Cell Report, Vol. 15, pp. 91-96 (1995)) describes genetically engineering resistance construct to potato virus X in four commercial potato cultivars. However, said document states that transgenic potato clones which have included the 3KG gene (the TGB3 construct). However, when these transgenic plants were challenged with PVX, there was no protection against PVX suggesting that the OK protein does not play a

role in the protection against PVX.

Aims of the invention

The present invention aims to provide a new
5 method for introducing various viral resistances into a cell and a plant and the viral resistant cell and plant obtained.

A main aim of the invention is to provide a new method for introducing BNYVV resistance into a cell and
10 a plant and the BNYVV-resistant cell and plant, in particular a sugar beet cell and plant (Beta vulgaris ssp.), obtained.

Summary of the invention

15 The present invention provides the use of an alternative sequence of plant virus, especially the BNYVV, to obtain a high degree of tolerance to the viral infection, in particular to ensure a rapid and total blocking of virus multiplication and diffusion mechanisms
20 in a plant, especially in the sugar beet plant (Beta vulgaris), including fodder beet, Swiss Whard and table beet, which may also be subject to this viral infection. Expression of the resistance will be obtained in transgenic cell and plant, especially sugar beet cells and plants
25 produced by the transformation method subject to the Patent Application WO95/10178 or by other transformation methods based on Agrobacterium tumefaciens or direct gene transfer. Because of its high efficiency, the transformation method as described in WO95/10178 enables the production of large
30 numbers of transformed plants, especially sugar beet plants, and will be preferred to develop transgenic plants which may be analysed and characterized for their level of

viral resistance, especially BNYVV Resistance, including their field evaluation.

The genome of beet necrotic yellow vein furovirus (BNYVV) consists of five plus-sense RNAs, two of which (RNAs 1 and 2) encode functions essential for infection of all plants while the other three (RNAs 3, 4 and 5) are implicated in vector-mediated infection of sugar beet (Beta vulgaris) roots (1). Cell-to-cell movement of BNYVV is governed by a set of three successive, slightly overlapping viral genes on RNA 2 known as the triple gene block (TGB) (2), which encode, in order, the viral proteins P42, P13 and P15 (gene products are designated by their calculated M_r in kilodalton (3).

In the following description, the TGB genes and the corresponding proteins will be identified by the following terms : TGB1, TGB2, TGB3 or by their encoded viral protein number P42, P13 and P15. TGB counterparts are present in other furoviruses (4, 5), and in potex-, carla- and hordeiviruses (6).

In the table 1 are represented viruses having a TGB3 sequence, the molecular weight of TGB3 of said viruses, their host and references.

Table 1

Virus	Size of TGB3	Host	Reference
Apple stem pitting virus	8 kDa	apple	Jelkman, J. Gen. Virol. 75, 1535-1542 (1994)
Blueberry scorch virus	7 kDa	blueberry	Cavaleer et al., J. Gen. Virol. 75, 711-720 (1994)
Potato virus M	7 kDa	potato	Zavriev et al., J. Gen. Virol. 72, 9-14 (1991)
White clover mosaic virus	8 kDa	clover	Forster et al., Nucl. Acids Res. 16, 291-303 (1988)
Cymbidium mosaic virus	10 kDa	orchid	Neo et al., Plant Mol. Biol. 19, 1027-1029 (1992)
Barley stripe mosaic virus	17 kDa	barley	Gustafson et al., Nucl. Acids Res. 14, 3895-3909 (1986)
Potato mop top virus	21 kDa	potato	Scott et al., J. Gen. Virol. 75, 3561-3568 (1994)
Peanut clump virus	17 kDa	peanut	Hertzog et al., J. Gen. Virol. 75, 3147-3155 (1994)
Beet soil-borne virus	22 kDa	sugar beet	Koenig et al., Virology 216, 202-207 (1996)

The Inventors propose herewith a new method for providing resistance to plant viruses into a plant by

5 blocking virus multiplication and diffusion mechanisms into

said plant, especially into its root tissue. In order to demonstrate said resistance, the Inventors describe hereafter the effect of the overexpression of TGB sequences alone or in combination upon BNYVV multiplication and
5 diffusion mechanism in plants of C. quinoa which are also the hosts of the BNYVV virus and which could be more easily manipulated by the man skilled in the art.

The Inventors have also made experiments upon Beta macrocarpa. These results have shown that it will be
10 possible to obtain also the transformation of plants by the method according to the invention and obtain expression of TGB3 gene by said plants. Therefore, as explained in the following description, said method could be used to obtain various viral resistances into various plants species
15 subject to infection by viruses characterized by the presence of a TGB3 sequence in their genome.

It is known that BNYVV does not require synthesis of viral coat protein for production of local lesions on leaves of hosts such as Chenopodium quinoa (7),
20 indicating that virion formation is not required for cell-to-cell movement.

However, the manner in which the TGB components assist in the movement process is not understood although computer-assisted sequence comparisons have
25 detected characteristic conserved sequences which may provide clues to their function. Thus, the 5'-proximal TGB protein (TGB1) invariably contains a series of sequence motifs characteristic of an ATP/GTP-binding helicase while the second protein (TGB2) always has two potentially
30 membrane-spanning hydrophobic domains separated by a hydrophilic sequence which contains a highly conserved peptide motif of unknown significance. The sequence and

size of the third TGB protein (TGB3) is more variable although the N-terminal portion is generally rather hydrophobic. Subgenomic RNAs with 5' termini mapping upstream of the BNYVV TGB1 and TGB2 open reading frames (ORFs) have been detected (Figure 1) but no such species has been reported for TGB3 of BNYVV (2), or of any other of the TGB-containing viruses. In the case of potato virus X (PVX; ref. 8) and barley stripe mosaic virus (BSMV; ref. 9), there is evidence that the TGB2 and TGB3 products are expressed from the same subgenomic RNA.

So far, no example has been reported of a virus in which the three TGB members are arranged differently on the same RNA or are parcelled out to different genome RNAs, suggesting that their association in a particular order might be important in regulating their function.

The present invention concerns a method for inducing viral resistance to a virus comprising a triple gene block (TGB) with the proviso that it is not the potato virus X. Said virus is preferably selected from the group consisting of the apple stem pitting virus, the blueberry scorch virus, the potato virus M, the white clover mosaic virus, the Cymbidium mosaic virus, the barley stripe mosaic virus, the potato mop top virus, the peanut clump virus and the beet soil-borne virus; said method comprises the following steps :

- preparing a nucleic acid construct comprising a nucleic acid sequence corresponding to at least 20% of the nucleic acid sequence of TGB3 of said virus or its corresponding cDNA, being operably linked to one or more regulatory sequence(s) active in a plant.

- transforming a plant cell with the nucleic acid construct, and possibly
- regenerating the transgenic plant from the transformed plant cell.

5 Preferably, the plant is a plant which may be infected by the above-described virus and is preferably selected from the group consisting of apple, blueberry, potato, clover, orchid, barley, peanut and sugar beet.

 The present invention concerns also the
10 obtained plant cell and transgenic (or transformed) plant (made of said plant cells) resistant to said viruses and comprising said nucleic acid construct.

 The Inventors have also discovered unexpectedly that it is possible to induce BNYSV-resistance
15 into a plant by a method which comprises the following steps :

- preparing a nucleic acid construct comprising a nucleic acid sequence corresponding to at least 70%, preferably at least 90%, of the nucleic acid sequence of comprised
20 between the nucleotides 3627 and 4025 of the 5' strand of the genomic or subgenomic RNA 2 of the BNYSV or its corresponding cDNA, being operably linked to one or more regulatory sequence(s) active in a plant,
- transforming a plant cell with said construct, and
25 possibly
- regenerating a transgenic plant from the transformed plant cell.

 The nucleic acid sequence comprised between the nucleotides 3627 and 4025 of the 5' strand of the
30 genomic or subgenomic RNA 2 encoding the P19 protein is described in the figure 6 and in the publication (3). Said nucleic acid sequence and the corresponding amino acid

sequence are described in the following specification as SEQ ID NO. 1.

Another aspect of the present invention concerns a plant cell and a transgenic plant (made of said
5 plant cells) resistant to BNYVV and comprising a nucleic acid construct having a nucleic acid sequence corresponding to at least 70%, preferably at least 90%, to the nucleic acid sequence comprised between the nucleotides 3627 and 4025 of the 5' strand of the genomic or subgenomic RNA 2 of
10 BNYVV or the corresponding cDNA, being operably linked to one or more regulatory sequence(s) active in the plant.

Preferably, said plant cell or transgenic plant (made of said plant cells) resistant to BNYVV is obtained by the method according to the invention.

15 The variants of the nucleic acid sequence described as SEQ ID NO. 1 comprise insertion, substitution or deletion of nucleotides encoding the same or different amino acid(s). Therefore, the present invention concerns also said variants of the nucleic acid sequence SEQ ID NO.
20 1 which present more than 70% homology with said nucleic acid sequence and which are preferably able to hybridize to said nucleic acid sequence in stringent or non-stringent conditions.

Preferably, said sequences are also able to
25 induce BNYVV resistance into a plant.

The terms "induce a viral resistance into a plant" mean inducing a possible reduction or a
30 significative delay into the appearance of infection symptoms, virus multiplication or its diffusion mechanisms into the plant, especially in the root tissue.

The regulatory sequence(s) of said nucleic acid sequence are promoter sequence(s) and terminator

sequence(s) active into a plant.

The nucleic acid construct may also include a selectable marker gene, which could be used to identify the transformed cell or plant and express the nucleic acid
5 construct according to the invention.

Preferably, the cell is a stomatal cell and the plant is a sugar beet (Beta vulgaris ssp.) made of said cells.

According to the invention, the promoter
10 sequence is a constitutive or foreigner vegetal promoter sequence, preferably selected from the group consisting of 35S Cauliflower Mosaic Virus promoter sequence, polyubiquitin Arabidopsis thaliana promoter (43), a promoter which is mainly active in root tissues such as the
15 par promoter of the haemoglobin gene from Perosponia andersonii (Landsman et al. Mol. Gen. Genet. 214 : 68-73 (1988)) or a mixture thereof.

A last aspect of the present invention is related to a transgenic plant tissue such as fruit, stem,
20 root, tuber, seed of the transgenic plant according to the invention or a reproducible structure (preferably selected from the group consisting of calluses, buds or embryos) obtained from the transgenic plant or the cell according to the invention.

25 The techniques of plant transformation, tissue culture and regeneration used in the method according to the invention are the ones well known by the person skilled in the art. Such techniques are preferably the ones described in the International Patent Applications
30 WO95/10178 or WO91/13159 corresponding to the European Patent Application EP-B-0517833, which are incorporated herein by reference. These techniques are preferably used

for the preparation of transgenic sugar beets according to the invention.

Brief description of the drawings

- 5 The figure 1 represents Structure of wild-type BNYVV RNA 2 and of replicons expressing TGB proteins.
- The figure 2 represents the In vitro translation of the replicons of the TGB genes in wheat germ extract.
- 10 The figure 3 represents the amplification of the replicons encoding the TGB proteins in *Chenopodium quinoa* protoplasts and expression of P42.
- The figure 4 represents the complementation of RNA 2 transcripts containing defects in the
- 15 different TGB genes by the corresponding wild-type gene supplied from a replicon.
- The figure 5 shows the effect of replicons on infection with wild-type BNYVV RNAs 1 and 2.
- The figure 6 represents the nucleotides and amino acid
- 20 sequence of TGB3 encoding the P15 of BNYVV.
- The figure 7 shows the presence of the coding regions of the BNYVV p15 gene in the sugar beet genome by a PCR.
- The figure 8 shows the integration of the BNYVV P15 gene
- 25 in the sugar beet genome by Southern blot hybridisation.

Description of the invention

In order to identify the potential use of

30 particular gene sequences isolated from the viral RNA2 of BNYVV and to create BNYVV resistance in a sugar beet via plant transformation, the inventors have investigated

whether independent expressions of the BNYVV TGB protein is possible by inserting the ORF of each into a viral replication-dependant "replicon" derived from BNYVV RNA3. The Inventors have showed that in mixed infections of C. quinoa leaves, the BNYVV TGB1 or TGB2 protein so expressed can complement BNYVV RNA 2 containing a mutation disabling the corresponding TGB protein. No complementation was observed with a replicon containing TGB3, however, unless the TGB3 ORF was positioned downstream of the ORF for TGB2 in the replicon. When co-inoculated with wild-type RNAs 1 and 2, the replicon expressing the TGB3 ORF of BNYVV inhibited infection. The data are consistent with a model for expression of the TGB proteins in which translation of P15 from a dicistronic subgenomic RNA regulates P15 expression levels in vivo. The Inventors have also identified that high expression of P15 could ensure a rapid and total blocking of virus multiplication and diffusion mechanisms in the plant.

20 MATERIALS AND METHODS

cDNA clones

The transcription vector for production of wild-type full-length BNYVV RNA 1 and RNA 2 were pB15 (10) and pB2-14 (11), respectively. Transcription vectors for previously described RNA 2 mutants were pB2-14-F, -H, -I and -J (2) and pB2-14- Δ SN, - Δ S12, - Δ S37, - Δ B1, - Δ B2, - Δ B2, - Δ N and -GAA (11). The RNA 2 deletion mutant pB2-14- HP1 was produced by elimination of the sequence between nucleotides 3158 and 3258. The empty BNYVV RNA 3-derived replicon, rep0, was obtained by transcription of the RNA 3 deletion mutant pB35AAES (12). TGB sequences for insertion into rep0 were amplified by the polymerase chain reaction

PCR using primers which each contained a non-templated BamHI site at their 5'-extremity. PCR fragments corresponding to the P42 gene (nucleotides 2127-3297), the P13 gene (nucleotides 3282-3650), the P15 gene (nucleotides 3627-4025) and both the P13 and P15 genes (nucleotides 3282-4025) were digested with BamHI and inserted into BamHI-cleaved pB35AΔES. The resulting constructs were used to transcribe rep42, rep13, rep15, and rep1315, respectively. A replicon containing a frameshift mutation in the P15 ORF (Rep15-X) was produced by filling in the overhangs of an insert XbaI site (nucleotide 3948). The insert frameshift mutations in rep13-I, rep1315-I and rep15-J were created as described for the corresponding mutations in full-length RNA 2 (2). Cloned PCR-amplified sequences were verified to be error-free by sequencing (13).

In vitro transcripts

Capped transcripts were prepared by bacteriophage T7 polymerase run-off transcription (10) of plasmid DNA linearized by HindIII for pB15 and the replicon constructs and by SalI for pB2-14 and related constructs. Transcript concentration and integrity were evaluated by agarose gel electrophoresis. Leaves were mechanically inoculated with 50 μ l per leaf of inoculation buffer containing 1 μ g of each transcript (21). In some experiments, the RNA 1 and 2 transcripts were replaced by 0.025 μ g of the highly infectious viral RNA purified from BNYVV isolate Stras 11-10. Preliminary experiments showed that this amount of viral RNA was approximately equivalent in infectivity as measured by a local lesion assay to a mixture containing 1 μ g each of the RNA 1 and 2

transcripts. For protoplast infections, 0.5 μ g of viral RNAs 1 and 2 plus 3 μ g replicon transcript were inoculated to $2 \cdot 10^5$ protoplasts by electroporation (2).

Transcripts obtained from replicons were translated in a wheat germ extract (14) and the [35 S]-labelled translation products were visualized by autoradiography after SDS-PAGE (15, 16). Radioactivity incorporated into translation products was quantified with a Fujix MAS1000 BioAnalyzer and the values were adjusted for methionine content in calculating relative translation levels.

Detection of viral RNA and proteins

Total RNA was extracted (2) from inoculated leaves 10 days post-inoculation (pi) and from protoplasts 48 hr pi. Viral RNA was detected by northern hybridization with 32 P-labelled antisense viral RNA transcripts (17) as probes. The RNA 1-specific probe was complementary to nucleotides 4740-5650, the RNA 2-specific probe to nucleotides 2324-3789 and the RNA 3-specific probe to nucleotides 1-380. P42, P14 and coat protein were detected by Western blot of total protein extracts of infected protoplasts using a rabbit polyclonal antiserum specific for each protein (18). The stability of mutations introduced into RNA 2 was tested by the polynucleotide chain reaction following reverse transcription (RT-PCR) of total RNA extracts from infected plants. Reverse transcripts were produced with an ExpandTM reverse transcription kit (Boehringer) following the manufacturer's instructions. PCR was carried out essentially as described (19) using 25 cycles of the following regimen: 94 °C 30

sec, 50 °C (30 sec), 72 °C (3 min). Primer pairs for PCR amplification of different regions of the RNA 3 cDNA corresponded to (or were complementary to, in the case of the second member of each pair of primers) nucleotides 1143-1151 and 3393-3412 (P42 gene), and nucleotides 3151-3169 and 4128-4148 (P13 and P15 genes). The primer used to initiate cDNA synthesis prior to the PCR reactions was complementary to nucleotides 4128-4148.

10 RESULTS

Replicons expressing the BNYVV TGB proteins

Provided that sufficient sequences at the 3'- and 5'-extremities are retained, a BNYVV RNA 3 transcript from which the central region has been deleted can replicate efficiently on *C. quinoa* leaves when coinoculated with RNAs 1 and 2, and can express a foreign gene inserted in place of the deleted sequence (12, 20). The Inventors have used such an RNA 3-based "replicon" to express each of the BNYVV TGB proteins out of their normal context in RNA 2 and tested the capacity of each replicon to complement an RNA 2 mutant defective in the corresponding TGB gene.

The replicons employed in this study are depicted in Figure 1. Figure 1 A is the Genome map of RNA 2. The TGB genes are shaded and lines above the map indicate the extents of the subgenomic RNAs 2sub₁ and 2sub₂. The positions of deletions and frameshift-inducing insertions in RNA 2 are indicated. The 5'-terminal cap structure is denoted by a circle. P21 is the major viral coat protein. RT - readthrough domain. B - BNYVV RNA 3-derived replicons containing the TGB genes of BNYVV light shading or the TGB₂ gene encoding P17 of peanut clump

virus (PCV) (dark shading). The *Bam*HI site in the empty replicon (rep0) used for insertion of the PCR-amplified TGB sequences is shown. The positions of frameshift-inducing insertions in the various P13 and P15 mutant replicons are indicated. In addition to the constructs rep42, rep13 and rep15, which each contain a TGB gene, a fourth construct (rep1315) was produced containing both the P13 and P15 genes arranged in the same relative configuration as in RNA 2. The ability of each replicon to direct expression of the inserted gene or genes was tested by in vitro translation of the transcript in a wheat germ extract. The rep42, rep13 and rep15 transcripts each directed synthesis of an abundant product (Figure 2A, lane 2; Figure 2B, lanes 2 and 3), which was not produced in translations programmed with transcript corresponding to the empty replicon, rep0 (Figure 2A, lane 1; Figure 2B, lane 1). In the figure 2 (A), are represented ³⁵S-methionine-labelled translation products of the empty replicon rep0 (lane 1) and rep42 (lane 2) displayed by autoradiography following PAGE (15). The indicated band was identified as P42 by comparison of its mobility to that of molecular weight markers (not shown). In the figure 2 (B), are represented translation products directed by rep0 (lane 1), rep13 (lane 2), rep15 (lane 3) and rep1315 (lane 4) displayed by autoradiography following PAGE (16). The bands tentatively identified as P13 and P15 are indicated to the right. The background band denoted by an asterisk was also synthesized when no transcript was introduced into the translation extract. The relative mobilities of the various translation products were as expected except that the putative P13 migrated slightly more slowly than P15, presumably because of its nontypical amino acid composition. The disjunctive

construct rep1315 directed synthesis of both P13 and P15 (Figure 2B, lane 4) in relative molar amounts of 3:1 (values corrected for the difference in methionine contents of the two proteins; if the N-terminal methionine of each protein is removed post-translationally, the molar ratio is 5:1).

The capacity of the replicons to be amplified by the viral replication machinery *in vivo* was tested by coinoculating replicon transcripts to *C. guinea* protoplasts along with BNYVV RNA's 1 and 2. Northern blot analysis of total RNA extracted from the protoplasts 48 hr pi revealed that all the replicons containing the TGB genes were efficiently amplified (Figure 3A). The figure 3 (A) represents detection by northern hybridization of viral RNAs in *C. guinea* protoplasts inoculated with BNYVV RNAs 1 and 2 alone (lane 2) or supplemented with rep0 (lane 3), rep42 (lane 4) rep13 (lane 5) rep1315 (lane 6) and rep15 (lane 7). RNA from mock-inoculated protoplasts was analyzed in lane 1. The protoplasts were harvested 48 hr pi and viral RNAs were detected using ³²P-labelled viral RNA-specific antisense RNA probes. The replicons are indicated by arrow heads. The figure 3 (B) represents Immunodetection of P42 in total protein extracts of *C. guinea* protoplasts inoculated with BNYVV RNAs 1 and 2 (lane 2), transcript of wild-type RNA 1 plus transcript of the RNA 2 mutant pB2-14-H, which contains a frameshift mutation in the P42 gene (lane 3), the RNA 1 and pB2-14-H transcripts plus rep42 (lane 4). Protein extracted from mock inoculated protoplasts was analyzed in lane 1. After PAGE and electrotransfer to nitrocellulose, P42, major viral coat protein CP and P14 were immunodetected with a mixture of antisera specific for each protein. The positions of

molecular weight standards are labelled in kilodaltons to the left of the blot. Western blot analysis revealed that the P42 level in protoplasts infected with a mixture of rep42 plus transcripts of RNA 1 and the frameshift mutant pB2-14-H, caused by filling in an *SpeI* site within the P42 gene of RNA 2 (see Figure 1), was about twice that in protoplasts infected with RNA 1 plus wild-type RNA 2 (Figure 3B, lanes 2 and 3). Note that the levels of accumulation of two other immunodetectable RNA 2 gene products (the major viral coat protein and P14; Figure 1) were not modified by the presence of rep42. P13 and P15 could not be immuno-detected in such experiments.

The BNYPV TGB proteins can be complemented in trans

The ability of the replicons containing the TGB genes to supply movement functions in whole plants was tested by coinoculating leaves of the local lesion host *C. quinoa* with one of a series of RNA 2 transcripts containing a mutation disabling a TGB gene plus a replicon containing the corresponding wild-type gene. In all experiments, the inoculum also contained transcript of wild-type RNA 1 as source of viral RNA-dependent RNA replicase, although this fact will not always be stated explicitly below. For the P42 gene, the RNA 2 mutants tested included the frameshift mutant (pB2-14-H) caused by filling in an *SpeI* site at nucleotide 2280, a series of mutants containing short in-frame deletions at different positions in the P42 ORF (mutants pB2-14-ΔS12, -ΔSN, -ΔB1, -ΔB2, -ΔN, and -ΔHP1; Figure 1A; also see ref. 11), and a deletion mutant (pB2-14-F; Figure 1) where removal of a 935 nucleotide sequence upstream of the P42 ORF has inactivated the promoter for the subgenomic RNA (RNA 2sub_g) responsible for P42

synthesis. Inocula containing RNA 1 transcript plus any one of the above mutant RNA 2 transcripts did not produce local lesions on *C. quinquefolia* and no progeny viral RNA could be detected in the inoculated leaves 10 days pi (Figure 4, lanes 3 and 5; see ref. 11 for the other mutants). In the figure 4, the replicon indicated at the top of each lane was inoculated to leaves of *C. quinquefolia* together with wild-type RNA 1 transcript plus either wild-type RNA 2 transcript (lane 2) or the mutant RNA 2 transcript identified above each lane. In lanes 19 and 20 the inoculum contained rep42 and rep15 (lane 19) or rep42 and rep1315 (lane 20), in addition to RNA 1 and pB2-14-HP1 transcripts. Lane 1 contains RNA from a non-inoculated control plant. Inoculated leaves were harvested 10 days pi and tested for viral RNA contents by northern hybridization as described in Figure 3. The positions of replicons are indicated by arrows. When rep42 transcript was included in the inoculum, numerous local lesions (20-80 per leaf) appeared on the inoculated leaves except for the inoculum containing transcript of the RNA 2 deletion mutant pB2-14-HP1, which remained symptomless. The resulting pale green lesions were similar in appearance to those elicited by inoculation with RNA 1 plus wild-type RNA 2 except for the RNA 2 mutant pB2-14-F, where necrotic local lesions were formed. In this latter case, the necrotic lesion phenotype may be related to production of a truncated form of the readthrough RT^h protein by this RNA 2 mutant (7).

Northern hybridization of the inoculated leaves 10 days pi revealed the presence of progeny viral RNAs of the length expected for RNAs 1, 2 and rep42 for all the RNA 2 mutants (Figure 4, lanes 2-4, 7-11) except the deletion mutant pB2-14-HP1 (Figure 4, lane 19). As will be

shown below, the failure of pB2-14-AHP1 to be complemented by rep42 is probably due to deletion of the promoter for the subgenomic RNA (RNA 2sub_B), which is believed to direct translation of the downstream TGB proteins.

5 Similar complementation experiments were carried out with rep13, rep15 and the dicistronic construct rep1315. Both rep13 and rep1315 were able to complement accumulation on leaves (Fig 4, lanes 14 and 15) of the mutant pB2-14-I, in which the P13 gene had been disabled by
10 insertion of four nucleotides (the insertion created an XhoI site), although the resulting local lesions were necrotic. Necrotic local lesions were also produced during mixed infections with the aforesaid replicons and wild-type RNA 1 and 2 (see below), indicating that the replicon-
15 related symptom phenotype is dominant over the wild-type. The novel symptoms may be related to differences in the time course of synthesis or the level of accumulation of P13 when it is expressed from the replicon rather than full-length RNA 2.

20 In experiments such as those described above, it is important to demonstrate that the mutation originally introduced into the P42 or P13 gene on the RNA 2 transcript was still present in the progeny RNA 2, that is, the defective copy of the TGB gene on the transcript had not
25 been converted to the wild-type by RNA recombination in planta (21) with the copy present on the replicon. Therefore, an RT-PCR experiment was carried out on the progeny viral RNA from a plant infected with RNA 1, pB2-14-I transcript (P42 gene disrupted by filling in an SpeI
30 site) and rep42. The primer pair used in the RT-PCR hybridized to RNA 2 sequences flanking the P42 gene and hence amplifies the copy of the gene present in RNA 2 but

not the copy on the replicon, where the flanking sequences are absent. Restriction enzyme analysis revealed that the *SpeI* site was absent in the resulting amplified DNA fragment, as expected for the mutated rather than the wild-type form of the TGB gene. Similar analysis of progeny viral RNA from plants infected with RNA 1, pB2-14-I (frameshift mutation in P13 gene creating an *XhoI* site, and either rep13 or rep1315 similarly demonstrated that the mutation disabling the copy of the P13 gene on the RNA 2 transcript was conserved in the progeny RNA 2. We conclude that rep42 and rep13 are indeed complementing P42 and P13 function by supplying the gene product in trans rather than simply serving as a source of the wild-type TGB sequence for recombination.

Unexpectedly, the replicon expressing the wild-type P15 gene (rep15) was unable to complement the P15-defective RNA 2 mutant pB2-14-J in mixed inoculations. No local lesions formed on the inoculated leaves 10 days pi and no viral RNA could be detected in the leaves by northern blot (Fig 4, lane 17). On the other hand, when pB2-14-J transcript was coinoculated with rep1315, local lesions (of the necrotic type) appeared and progeny viral RNAs were readily detected (Figure 4, lane 18). In this latter case, analysis of an RT-PCR product containing the P15 gene in the progeny RNA 2 revealed that the mutation disabling the gene was still present. Complementation of pB2-14-J still occurred when the P13 ORF in the dicistronic replicon was interrupted by a frameshift mutation (rep1315-1; Figure 1) establishing that expression of full-length P13 from the first ORF of the dicistronic replicon is not required for complementation by the downstream copy of the P15 gene.

Evidence that P15 is expressed from a dicistronic subgenomic RNA

An RNA 2-derived subgenomic RNA (RNA 2sub_b) of about 1500 nucleotides length has been detected in
5 BNYVV-infected tissue (2). The 5'-extremity of this species has not been mapped precisely but is predicted to lie near the 5'-terminus of the P13 ORF. No subgenomic RNA with 5'-end upstream of the P15 ORF has been detected, raising the possibility that, as in BSMV (8), P13 and P15 are both
10 expressed from RNA 2sub_b.

The aforementioned inability of rep42 to complement the P42-defective RNA 2 mutant pB2-14- HP1 could stem from polar effects of the RNA 2 deletion on synthesis of downstream TGB proteins if the deletion has disabled the
15 RNA 2sub_b promoter (The right-hand boundary of the deletion in pB2-14-ΔHP1 is only 30 residues upstream of the P13 initiation codon). To test this hypothesis, an experiment was carried out in which the pB2-14-ΔHP1 transcript was complemented with both rep42 and rep1315. Leaves inoculated
20 with this mixture developed local lesions and contained progeny viral RNA's (Figure 4, lane 20). If, on the other hand, rep13 rather than rep1315 was used along with rep42 to complement pB2-24- HP1, no symptoms appeared and no progeny viral RNA's were detected by northern blot (Figure
25 4, lane 19). These observations are consistent with the hypothesis that the pB2-14- HP1 deletion interferes with expression of the downstream TGB ORF's, presumably by blocking RNA 2sub_b transcription. Furthermore, the fact that complementation was successful with rep1315 but not
30 with rep13 indicates that P15 as well as P13 is translated from RNA 2sub_b.

Independent expression of P15 inhibits infection with wild-type viral RNA

The ability of rep1315 but not rep15 to complement the P15-defective RNA 2 mutant pB2-14-J in leaf infections could indicate that independent expression of P15 from the monocistronic replicon interferes with the viral infection cycle by producing the gene product in excessive quantities relative to P13. To test this hypothesis, an experiment was carried out in which rep15 was inoculated to C. Quinoa leaves along with wild-type viral RNAs 1 and 2. No lesions appeared on the inoculated leaves, even at late times pi (Figure 5A), and no viral RNA could be detected by northern blot (Figure 5B, lane 6). The figure 5 (A) represents leaves of C. quinoa inoculated with RNAs 1 and 2 (left) or RNAs 1 and 2 plus rep15 (right). The leaves were photographed 20 days pi when the local lesions on the leaf to the left had expanded so as to cover much of the leaf surface. In the figure 5 (B), analysis by northern hybridization (as described in Figure 3) of the viral RNA contents of C. quinoa leaves inoculated with BNYVV RNAs 1 and 2 alone (lane 1) or together with rep0 (lane 2), rep42 (lane 3), rep13 (lane 4), rep1315 (lane 5), rep15 (lane 6), rep15-J (lane 7), rep15-X (lane 8) or repPCV-P17 (lane 9). The positions of replicons are indicated by arrows. (C) Analysis by northern hybridization of the viral RNA contents of the inoculated leaves (lanes 1, 3 and 5) and the roots (lanes 2, 4 and 6) of Beta macrocarpa either mock-inoculated (lanes 1 and 2), inoculated with BNYVV RNAs 1, 2 and 3 (lanes 3 and 4) or with RNAs 1, 2 and 3 plus rep15 (lanes 5 and 6). RNA 3 was included in the inoculum because it is necessary for systemic movement in B. macrocarpa (11). Under these conditions, leaves inoculated

with RNAs 1 and 2 alone were heavily infected (Figure 5A; Figure 5B, lane 2). The inhibition of virus infection by repl5 was dose-dependent. Addition of ten times less repl5 to the inoculum mix still resulted in almost complete inhibition of lesion formation but lesser amounts of the replicon were progressively less effective in blocking the infection. Repl5 also blocked the appearance of progeny viral RNA in the inoculated leaves and in the roots of Beta macrocarpa, a systemic host of BNYVV (Figure 5C, lanes 5 and 6). The empty replicon, rep0, and replicons expressing the other two TGB proteins (rep42, repl3, repl315), on the other hand, did not significantly inhibit BNYVV infection of C. guinea leaves (Figure 5B, lanes 2-5).

Since repl5 did not interfere with amplification of RNA 1 and 2 in C. guinea protoplasts (see Figure 3), this suggests that the replicon interferes with movement of the virus from the initial site of infection into neighbouring cells (cell-to-cell movement) during local lesion formation on leaves. Lesion formation was not inhibited by coinoculation of Stras 12 RNA with the replicons repl5-J or repl5-X (Figure 5B, lanes 7 and 8), which encode frameshift-truncated forms of P15. This finding confirms that expression of P15 from the replicon, rather than the simple presence of the corresponding RNA sequence, is required for inhibition during the mixed infection experiments. In the presence of repl5-X, however, the resulting local lesions were about one third the diameter of the lesions formed by infection with Stras 12 alone or with Stras 12 plus repl5-J and the content of progeny viral RNA in the infected leaves was significantly lower (Figure 5B, lane 8). This finding suggests that the almost full length P15 molecule produced by repl5-X,

although incapable of substituting for wild-type P15 in a complementation experiment can interfere with the cell-to-cell movement activity of the wild-type P15 produced from RNA 2. Presumably, the full-length and truncated forms of P15 compete with one another for binding sites on another component (which could be of either viral or cellular origin) involved in the movement process.

As noted above, sequence comparisons between different viruses possessing a TGB have revealed little sequence similarity between the different TGB3 genes. For example, the 17 kDa TGB3 protein (P17) of peanut clump virus (PCV) displays no significant sequence similarity with P15 of BNYSV (4) even though both viruses can infect C. guinea. To determine whether independent expression of the PCV TGB3 protein can interfere with a BNYSV infection in a manner similar to that observed with rep15, a BNYSV RNA 3-derived replicon containing the PCV TGB3 (repPCV-P17; Figure 1) was constructed. C. guinea leaves inoculated with BNYSV RNAs 1 and 2 plus repPCV-P17 did not develop symptoms and no progeny BNYSV RNA's could be detected by northern blot (Figure 5A, lane 9). This observation suggests that the pathways by which BNYSV and PCV move from cell-to-cell in C. guinea share at least one common element which in spite of their dissimilarity in sequence, interacts with the TGB3 products of both viruses.

The Inventors have shown that replicons carrying P42 and P13 can complement BNYSV RNA 2 carrying the corresponding defective gene but that a replicon carrying P15 cannot. In the latter case, complementation can occur, however, if the P15 gene is supplied as the second gene on a dicistronic RNA (rep1315) carrying the P13 gene in first position. It should be noted that the

relative disposition of the P13 and P15 genes on rep1315 is identical to their disposition on RNA 2sub₂, the subgenomic RNA which is believed to direct synthesis of both proteins in wild-type infections. This suggests that successful

5 cell-to-cell movement of BNYVV requires the presence of P13 and P15 in appropriate relative amounts and that production of both proteins from the same subgenomic RNA represents a mechanism for coordinating their synthesis. The inability of rep15 to complement the P15 mutant RNA 2 transcript pB2-

10 14-J and its ability to inhibit infection by wild-type virus would then both be due to over-production of P15 relative to P13 when the former is translated from the replicon and the latter from RNA 2. When P15 is expressed from the dicistronic replicon rep1315, on the other hand,

15 appropriate relative P13-P15 levels would be produced, allowing cell-to-cell movement to proceed. The "correct" relative levels of accumulation of P13 and P15 in a wild-type infection are not known. Translation of rep1315 in wheat germ extract produced three to five times more P13

20 than P15 but such experiments do not necessarily reflect the situation in planta since the turnover rates of the two proteins may differ significantly. Note that these results indicate that TGB-mediated cell-to-cell movement is less sensitive to over-expression of P42 and P13 relative to the

25 "correct" levels characteristic of a normal infection since co-inoculation of rep42, rep13 or rep1315 with wild-type virus did not inhibit infection (Figure 5, lanes 3-5), although the lesions produced in the presence of rep13 and rep1315 were necrotic. Thus, it shows that expression of

30 P15 in transgenic plants could provide a mechanism for inducing BNYVV-resistance ("pathogen-derived resistance"; ref. 23) in such plants, provided that sufficient P15

expression levels can be attained.

To gain a better understanding of how the relative levels of P13 and P15 are regulated during translation will require learning how the P15 cistron is accessed by ribosomes on RNA 2sub₀. Translation initiation at an internal cistron of a eucaryotic messenger RNA may occur by several mechanisms, including (i) leaky scanning, where a fraction of the ribosomal subunits which begin scanning the RNA at the 5'-end move past the first (non-optimal) upstream AUG without initiating (24), and (ii) internal entry, where ribosomal subunits bind directly to a special sequence on the RNA near the internal initiation codon (25). A third possible mechanism, termination-reinitiation (24), appears unlikely to apply to any of the TGB-containing viruses because the overlap between the TGB2 and TGB3 cistrons would require ribosomes to scan backwards after terminating TGB2 to reach the TGB3 initiation codon. It has been suggested that the TGB3 proteins of BSMV and PVX are translated by a leaky scanning mechanism (8, 9). The BNYVV P15 gene may also be produced by leaky scanning although it should be noted, however, that the context of the BNYVV P13 initiation codon (AUAAUGU) is nearly optimal and there are also two downstream AUG's which scanning subunits would have to ignore to reach the P15 initiation codon.

EXAMPLES

The following examples are transformation of plant made by the technique described in the International Patent Application WO95 11174 incorporated herein by reference.

The plant material and growth conditions were the ones described by Hall et al., Plant Cell Reports 12, pp. 339-342 (1993) Pedersen et al., Plant Science 95, pp. 89-97 (1993), and Hall et al, Nature biotechnology 14, 5 1996, in press.

Plasmid vectors and DNA preparation

The plasmid pET-P15 (harbouring the P15 nucleic acid sequence) was restricted at its single BamHI 10 site and blunt-ended with T4 DNA polymerase. After purification by electrophoresis in 0.8% agarose gel, the linear plasmid was restricted at its single NcoI site. The P15 gene fragment of 400 bp was purified by electrophoresis and inserted into pMJBX-Ub (harbouring the Arabidopsis 15 polyubiquitin promoter (Norris et al., Plant Molecular Biology 21, pp. 895-906 (1993), a TMV enhancer sequence and the Nos 3' terminator) cut with NcoI and SmaI restriction endonucleases. In the plasmid so obtained (pMJBX-Ub-P15), the nucleic acid sequence of the P15 gene 20 is placed under the control of the Arabidopsis polyubiquitin promoter followed by the TMV enhancer sequence. The EcoRI fragment from plasmid pB235Sack contains the pat gene, used as the selective marker, encoding phosphinothricin acetyl transferase (obtained from 25 Agrevo, Berlin Germany). On this EcoRI fragment, the nucleic acid sequence of the pat gene is under the control of the 5' and 3' expression signals of the Cauliflower virus. The plasmid pMJBS6, resulting from the combination of this EcoRI-pat fragment and a partial EcoRI digestion of 30 plasmid pMJBX-Ub-P15, contains both the pat and the P15 genes. This pMJBS6 plasmid is a high-copy plasmid based on the pUC18 vector and contains also the tetracycline resistance gene

amp^r. In the plasmid pIGPD7, harbouring the same pat
fragment as pB235SAck, the β -lactamase gene was replaced by
an *igpd* (imidazole glycerol phosphate dehydratase) gene
from *Saccharomyces cerevisiae* (Struhl et al., Proceedings
5 of the National Academy of Science USA 73, pp. 1471-1475
(1976)). Selection for and maintenance of the plasmid in
Escherichia coli was achieved by complementation of an
auxotrophic *hisB* strain SB3930 on minimal medium in the
absence of antibiotics. The P15 fragment, with its
10 ubiquitin promoter and terminator sequence, was purified as
a 2500 bp fragment obtained from the pMJBX-Ub-P15 plasmid
after it was cut at the single HindIII site, followed by a
partial EcoRI restriction. This fragment was blunt-ended
and inserted in a blunt-ended pIGPD7 plasmid, cut at the
15 single NcoI site. The resulting pIGPDS4 plasmid contains
both the *pat* and the P15 genes on a vector without the β -
lactamase gene.

Plant material

20 In vitro shoot cultures of sugar beet
plantlets were initiated to provide a reusable and uniform
source of sterile starting material and were maintained
with a 4-weekly subculture period as described by Hall et
al., Plant Cell Reports 12, pp. 339-342 (1993).

25

Large-scale isolation of sugar beet epidermis

A modified version of the blender method by
Kruse et al., Plant Physiology 690, pp. 1383-1384 (1989)
was used. For each isolation, 2 g leaves with the midribs
30 removed from 4 week old shoots was blended in a Waring
blender at maximum speed (1200 rpm) for 60 sec in a 250 ml

metal beaker containing 50 ml cold (4 °C) Ficoll medium (100 g /l Ficoll, 735 mg/l $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 1 g/l PVP40, autoclaved). The epidermal fragments were then recovered on a 297 μm nylon filter and washed with 500 ml sterile tap
5 water. These were rinsed from the filter into a 9 cm Petri dish using 10 ml CPW9M containing 3.8% (w/v) $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ (Krens et al., Theoretical and Applied Genetics 79, pp. 390-396 (1990)). Any remaining leaf fragments were removed and dishes were preincubated for 1 h at room temperature.

10

Guard cell protoplast isolation from enriched epidermis fractions

To recover the epidermis fraction, the suspension was centrifuged for 1 min at 55 x g after which
15 the supernatant was removed. The pellet was resuspended in 50 ml enzyme mix and 5 ml aliquots were transferred to each of 10, 6 cm Petri dishes (Greiner, TC quality), sealed with parafilm and incubated overnight at 25 °C in darkness with gentle agitation. The digestion medium consisted of CPW9M
20 supplemented with 0.5% (w/v) Cellulase RS and 3% (w/v) Macerozyme R10 (Yakult Honsha, Tokyo, Japan), pH 5.8. The following morning, the protoplasts were generally seen floating near the surface of the digestion mix. After gentle agitation of the suspensions using a sterile pipette
25 to release the protoplasts still adhering to cuticle fragments, the digestions were pooled and passed through 297 and 55 μm nylon filters. The filtrate was mixed with an equal volume of iso-osmotic Percoll containing 15% (w/v) sucrose (Percoll15S) and divided over 12 x 12 ml centrifuge
30 tubes. In each tube, first 1 ml CPW15S (Krens et al., 1990) and then 0.5 ml 9% (w/v) mannitol containing 1 mM CaCl_2 .

(9M) were carefully layered on top of the protoplast suspension. After centrifugation at 55 x g for 10 min the viable guard cells were visible in bands at the CPW16S/9M interface. To concentrate the protoplasts, these bands were
5 collected and mixed with Percoll15S to give a final volume of 16 ml. This was then divided between 2 centrifuge tubes, layered as above and recentrifuged. Careful removal of the 9M layers yielded the enriched guard cell protoplast fraction for subsequent counting using a haemocytometer.

10

Protoplast transformation

Transformations were performed in 12 ml centrifuge tubes, each containing 1×10^6 protoplasts suspended in 0.75 ml 9M medium. Plasmid DNA (50 μ g of
15 pMJBX-Ub-P15 and pIGPDS4) was added and, immediately after mixing, 0.75 ml PEG medium was added dropwise (40% PEG 6000 dissolved in F medium (Krens et al., Nature 296, pp. 72-74, (1982)). After thorough mixing, the suspension was kept at room temperature for 30 min with intermittent agitation.
20 Subsequently, at 5 min intervals, 4 x 3 ml aliquots of F medium were added. After centrifugation for 5 min at 55 x g the supernatant was discarded and the protoplast pellet resuspended in 9M and recentrifuged. The cells were finally resuspended in 1 ml of 9M medium for counting.

25

Protoplast culture and selection

Protoplasts were embedded in Ca alginate and cultured in modified, liquid KSP medium (Hall et al., 1990). To select for stably transformed cells, bialaphos, the active compound of Herbiace (Meiji Seika Ltd, Japan
30 was added after 7 days, to give a final concentration of 200 μ g/l. On day 14, the alginate discs were cut into 5 mm

slices and transferred to PGo medium (De Greef and Jacobs, Plant Science Letters 17, pp. 55-61, 1979) supplemented with 1 μ M BAP (PG1B) and 250 μ g/l bialaphos and solidified with 0.8% agarose.

5

Callus culture and regeneration

After 21 days, the pieces of alginate containing the non-visible microcalli were transferred to 9 cm Petri dishes containing 20 ml Medium K (3% sucrose, 0.8% agarose, 1 μ M BAP, PGo medium, pH = 5.8 autoclaved).
10 Culture was in the dark as above.

Friable, watery-type calli on reaching the size of approximately 1-2 mm in diameter, were individually picked off and cultured in groups of 20 on fresh Medium K.
15 At this stage, PCR analyses confirmed the presence of transformants.

At two-weekly intervals all calli were subcultured on to fresh medium.

20 Regenerants appeared during the first 8 weeks of culture of individual calli. When the first shoots were visible and had reached a size of approximately 2 mm, the dish was transferred into the light (3000 lux), 25 $^{\circ}$ C, 15 hour day length.

25 Plantlets approximately 4 mm long were transferred to individual culture tubes containing 15 ml of Medium K and were further subcultured in the light as above.

30 Rooting and transfer to the soil

When the plantlets had reached the four-leaf stages (usually after 5 to 6 weeks with one subculture

after 3 weeks, they were transferred to culture tubes containing 15 ml of Medium L (3% sucrose, 0.8% agarose, 25 μ M indolebutyric acid IBA), PGo medium, pH = 5.8 autoclaved (PGo medium described by De Greef W. et al.,
5 Plants Science Letters 17, pp. 55-61 (1979)) and further cultured as above.

When at least one root had reached a length of 1 cm, the plantlets were removed from the culture tubes and washed under running tap water to remove all fragments
10 of the agar, and transferred to soil in 9 cm pots in the greenhouse.

Plantlets were covered with a transparent plastic cup to provide a humid environment for 7 days, after which they could be grown without protection.

15 The plant transformed by the sequence SEQ ID NO. 1 according to the invention is recovered and has been expressed P15.

DNA analysis

20 Genomic DNA isolated from the primary transformants is electrophoresed in a 0.8% agarose gel after treatment with restriction enzymes and transferred to a nitrocellulose membrane using standard procedures, according to the manufacturer's protocol. Hybridisation is
25 performed with the DNA, as α^{32} P-dATP-labelled probes, whose presence it is desired to establish. The membranes were washed to a final stringency of 0.1% x SSC, 0.1% SDS at 60 °C. The hybridized DNA is visualised by darkening the X-ray film for 24 to 48 hours.

PCR analysis

Standard PCR techniques were used to detect a range of intact plasmid sequences. Reactions were performed using 25 cycles of 1 min denaturation at 94 °C, 1 min annealing; 2 min extension at 72 °C, with a final extension period of 5 min. The annealing temperatures were optimized for each primer combination. The presence of the coding region of the BNYVV P15 gene in the sugarbeet genome was verified by PCR using a pair of oligonucleotides as primers

10 : MOV1, sense primer [5'-GGTGCTTGTGTTAAAGTAGATTTATC-3' (nucleotides 3 to 29 on SEQ ID NO 1)] and MOV2 antisense primer [5'-CTATGATACCAAAACCAAACTATAGAC-3' (complementary to the nucleotides 369 to 395 on SEQ ID NO 1)]. This 393 bp long fragment comprises the whole coding region of the P15

15 gene for BNYVV (see figure 7).

Figure 7: Analysis of PCR products obtained with sugarbeet DNA from P15-transformants (lane 5 to 7) and an untransformed plant (lane 4). Low DNA Mass Ladder® (Life Technologies) was used as a size marker (lane 1).

20 Lane 2 and 3 correspond to the positive controls (pMJBS6/pIGPDS4). The arrow on the left shows the position of the expected PCR product.

Southern blot hybridisation analysis

25 The integration of the BNYVV P15 gene in the sugarbeet genome was verified by Southern blot hybridisation. Total DNA of primary transgenic regenerants was isolated, digested with restriction enzymes (PstI, KpnI, NcoI, SacI), electrophoresed, blotted and hybridised

30 with BNYVV P15 specific ³²P-labelled probes using PCR amplified MOV1-MOV2 fragment (see figure 8).

Figure 3: Southern blot analysis. Lambda DNA digested with HindIII was used as a size marker (lane 1). Lanes 2 to 16, DNA of the transgenics: 2 to 4 digested with SacI, 6 to 8 digested with PstI, 10 to 12 digested with NcoI, 14 to 16 digested with KpnI. Lanes 5, 9, 13 and 17 correspond to the untransformed plant.

REFERENCES

1. Richards K.E. & Tamada T., *Annu. Rev. Phytopathol.* 30, pp. 291-313 (1992)
2. Gilmer D. et al., *Virology* 189, pp. 40-47 (1992)
- 5 3. Bouzoubaa S. et al., *J. Gen. Virol.* 68, pp. 615-626 (1987)
4. Herzog E. et al., *J. Gen. Virol.* 18, pp. 3147-3155 (1994)
5. Scott K. P. et al., *J. Gen. Virol.* 75, pp. 3561-3568 (1994)
- 10 6. Koonin E.V. & Dolja V.V., *Crit. Rev. Biochem. and Mol. Biol.* 28, pp. 375-430 (1993)
7. Schmitt C. et al., *Proc. Natl. Acad. Sci. USA.* 89, pp. 5715-5719 (1992)
- 15 8. Morozov S.Y. et al., *J. Gen. Virol.* 72, pp. 2039-2042 (1991)
9. Zhou H. & Jackson A.O., *Virology* 216, pp. 367-379 (1996)
10. Quillet L. et al., *Virology* 172, pp. 293-301 (1989)
- 20 11. Bleykasten et al., *J. Gen. Virol.* 77, pp. 889-897 (1996)
12. Jupin I. et al., *Virology* 178, pp. 273-280 (1990)
13. Sanger F. et al., *Proc. Natl. Acad. Sci. USA* 74, pp. 5463-5467 (1977)
- 25 14. Fritsch C. et al., *J. Gen. Virol.* 46, pp. 381-389 (1980)
15. Laemmli U.K., *Nature* 227, pp. 680-685 (1972)
16. Schagger H. & von Jagow G., *Anal. Biochem.* 166, pp. 368-379 (1987)
- 30 17. Lemaire O. et al., *Virology* 162, pp. 232-235 (1988)
18. Niesbach-Klösigen U. et al., *Virology* 178, pp. 52-61 (1990)

19. Hehn A. et al., *Virology* 210, pp. 73-81 (1995)
20. Tamada T. et al., *J. Gen. Virol.* 70, pp. 3399-3409
1989
21. Kozak M., *J. Cell. Biol.* 108, pp. 299-241 (1989)
- 5 22. Pelletier J. & Sonenberg N, *Nature* 334, pp. 320-325
(1988)
23. Tamada T. & Baba T., *Annals of the Phytopathological
Society of Japan* 39, pp. 325-332 (1973)
24. Kuszala M. & Putz G., *Annals of Phytopathology* 9,
10 pp. 435-446 (1977)
25. Heskin B., *Archiv für Mikrobiology* 49, pp. 343-374
(1964)
26. Asher M.J.C., *Rhizomania In The sugar beet crop*, ed.
D.A. Cooke and R.K. Scott, Chapman & Hall, London,
15 pp. 312-338 (1993)
27. Richard-Molard M., *Rhizomanie In Institut français de
la betterave industrielle. Compte-rendu des travaux
effectués en 1994*, ITB, Paris pp. 225-229 (1995)
28. Henry C.M. et al, *Plant Pathology* 41, pp. 483-489
20 (1992)
29. Grassi G. et al., *Phytopath. Medit.* 28, pp. 131-139
(1989)
30. Merdinoglu D. et al., *Acad. Agric. Fr.* 79, n° 6,
pp. 85-98 (1993)
- 25 31. Scolten O.E. et al., *Archives of Virology* 136, pp. 349-
361 (1994)
32. Büttner G. & Bückky K., *Proceedings of the First
Symposium of the International Working Group on Plant
Viruses with Fungal Vectors*, Braunschweig Germany,
30 August 21-24 1990
33. Whitney E.D., *Plant Disease* 73, pp. 287-289 1989

34. Powell A.P. et al., *Science* **232**, pp. 738-743 (1986)
35. Fritchen J.H. & Beachy R.N., *Ann. Rev. Microbiol.* **47**, pp. 739-763 (1993)
36. Wilson T.M.A., *Proc. Natl. Acad. Sci. USA* **90**, pp. 3134-3141 (1993)
- 5
37. Gonsalves D. & Slightom J.L., *Seminars in Virology* **4**, pp. 397-405 (1993)
38. D'Halluin K. et al., *Biotechnology* **10**, pp. 309-314 (1992)
- 10
39. Kallerhof J. et al., *Plant Cell Reports* **9**, pp. 224-228 (1990)
40. Ehlers U. et al., *Theoretical and Applied Genetic* **81**, pp. 777-782 (1991)
41. Kraus J. et al., *Field performance of transgenic sugar beet plants expresing BNYVV coat protein plants, Fourth International Congress of Plant Molecular Biology, Int. Soc. for Plant Molecular Biology, Amsterdam (1994)*
- 15
42. Maiss E. et al., *Proceedings of the Third International Symposium on the Biosafety Results of Field Tests of Genetically Modified Plants and Microorganisms, Monterey, pp. 129-139 (1994)*
- 20
43. Norris et al., *Plant Molecular Biology* **21**, pp. 895-906 (1993)

CLAIMS

1. Method for inducing resistance to a virus comprising a TGB3 sequence with the proviso that it is not the potato virus X, into a plant cell or a plant,
5 comprising the following steps :
- preparing a nucleic acid construct comprising a nucleic acid sequence corresponding to at least 70% of the nucleic acid sequence of TGB3 of said virus or its
10 corresponding cDNA, being operably linked to one or more regulatory sequence(s) active in a plant,
 - transforming a plant cell with the nucleic acid construct, and possibly
 - regenerating a transgenic plant from the transformed
15 plant cell.
2. Method according to the claim 1, characterized in that the nucleic acid sequence of the nucleic acid construct corresponds to at least 90% of the nucleic acid sequence of TGB3 of said virus or its
20 complementary cDNA.
3. Method according to the claim 1 or 2, characterized in that the virus is selected from the group consisting of the apple stem pitting virus, the blueberry scorch virus, the potato virus M, the white clover mosaic
25 virus, the *Cymbidium* mosaic virus, the barley stripe mosaic virus, the potato mop top virus, the peanut clump virus and the beet soil-borne virus.
4. Method according to any of the preceding claims, characterized in that the plant cell is a stomatal
30 cell.
5. Method according to any of the preceding claims characterized in that the plant is selected from

the group consisting of apple, blueberry, potato, clover, orchid, barley or peanut.

6. Method according to claim 1 or 2, characterized in that the virus is BNYVV, the nucleic acid
5 sequence of TGB3 of said virus is comprised between the nucleotide 3627 and 4025 of the 5' strand of genomic or subgenomic RNA 2 of the BNYVV and the plant is a beet, preferably a sugar beet (*Beta vulgaris*).

7. Method according to any of the preceding
10 claims, characterized in that the regulatory sequence comprises a promoter sequence or a terminator sequence active in a plant.

8. Method according to claim 7 characterized in that the promoter sequence is a constitutive or a
15 foreigner vegetal promoter sequence.

9. Method according to the preceding claim 7, characterized in that the promoter sequence is selected from the group consisting of 35S Cauliflower Mosaic Virus promoter, and/or the polyubiquitin *Arabidopsis thaliana*
20 promoter.

10. Method according to any of the claim 7 to 9, characterized in that the promoter sequence is a promoter which is mainly active in the root tissue of plants such as the par promoter of the haemoglobin gene
25 from *Perosponia andersonii*.

11. Transgenic plant resistant to a virus with the proviso that it is not the potato virus X, comprising a nucleic acid construct having a nucleic acid sequence corresponding to at least 70% of the nucleic acid
30 sequence of TGB3 of said virus or its corresponding cDNA, being operably linked to one or more regulatory sequence(s) active in a plant.

12. Transgenic plant according to the claim 11, characterized in that the nucleic acid construct has a nucleic acid sequence corresponding to at least 90% of the nucleic acid sequence of TGB3 of said virus or its
5 complementary cDNA.

13. Transgenic plant according to the claim 11 or 12, characterized in that the virus is selected from the group consisting of the apple stem pitting virus, the blueberry scorch virus, the potato virus M, the white
10 clover mosaic virus, the *Cymbidium* mosaic virus, the potato virus X, the barley stripe mosaic virus, the potato mop top virus, the peanut clump virus and the beet soil-borne virus.

14. Transgenic plant according to the claims 15 11 to 13 being a plant selected from the group consisting of apple, blueberry, potato, clover, orchid, barley or peanut.

15. Transgenic plant according to the claims 11 or 12, characterized in that the transgenic plant being
20 a beet, preferably a sugar beet (*Beta vulgaris*) the virus is BNYSV and the nucleic acid sequence of TGB3 of said virus is comprised between the nucleotide 3627 and 4025 of the 5' strand of genomic or subgenomic RNA 2 of BNYSV or its corresponding cDNA.

25 16. Transgenic plant according to any of the preceding claims 11 to 15, characterized in that the regulatory sequence comprises a promoter sequence and a terminator sequence active in a plant.

17. Transgenic plant according to any of the
30 preceding claims 11 to 16, characterized in that the regulatory sequence comprises a promoter sequence which is a constitutive or a foreigner vegetal promoter sequence.

18. Transgenic plant according to the claim
17, characterized in that promoter sequence is selected
from the group consisting of 35S Cauliflower Mosaic Virus
promoter, and/or the polyubiquitin *Arabidopsis thaliana*
5 promoter.

19. Transgenic plant according to claim 17 or
18 characterized in that the promoter sequence is a
promoter which is mainly active in root tissues such as the
par promoter of the haemoglobin gene from *Perosponia*
10 *andersonii*.

20. Transgenic plant tissue selected from the
group consisting of fruit, stem, root, tuber, seed of a
plant according to any of the preceding claims 11 to 19.

21. Reproducible structure obtained from a
15 transgenic plant according to any of the preceding claims
11 to 19.




FIG. 1




FIG. 2

2/5

FIG. 3




FIG. 4





FIG. 5


1/1 31/11
ATG TTG GTT GTG GTT AAA GTA GAT TTA TCT AAT ATT GTA TTG TAC ATA GTT GCG GGT TGT
M V L V V K V D L S N I V L Y I V A G C
61/11 91/31
GTT GTT GTG AGT ATG TTG TAC TCA CCG TTT TTC AGC AAC GAT GTT AAA GCG TCC AGC TAT
V V V S M L Y S P F P S N D V K A S S Y
121/41 151/51
GCG GGA GCA ATT TTT AAG GGG AGC GGC TGT ATC ATG GAC AGG AAT TCG TTT GGT CAA TTT
A G A I P K G S G C I M D R N S P A Q P
181/61 211/71
GGG AGT TGC GAT ATT CCA AAG CAT GTA GCG GAG TCC ATC ACT AAG GTT GCG ACC AAA GAG
G S C D I P K E V A E S I T K V A T K E
241/81 271/91
TAC GAT GTT GAC ATA ATG GTA AAA AGG GGT GAA GTG ACC GTT GGT GTT GTG ACT CTC ACC
E D V D I M V K R G E V T V R V V T L T
301/101 331/111
GAA ACT AAT TTT ATA ATA TTA TCT AGA TTG TTT GGT TTG GCG GTG TTT TTG TTC ATG ATA
E T I P I I L S R L F G L A V P L F M I
361/121 391/131
TGT TTA ATG TCT ATA GTT TGG TTT TGG TAT CAT AGA TAA
C L M S I V W F W Y E R *

FIG. 6SEQ ID NO. 1



FIG. 7

FIG. 8



INTERNATIONAL SEARCH REPORT

International Application No.

PCT/BE 97/00092

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/82 A01H5/00 //C07K14/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
A	H. XU ET AL.: "Genetically engineered resistance to potato virus X in four commercial potato cultivars" PLANT CELL REP., vol. 15, 1995, pages 91-96, XP000617900 see abstract and page 95.	1,11
A	WO 91 13159 A (BIOSEM) 5 September 1991 cited in the application see Examples 15-18 and claims.	1



Further documents are listed in the continuation of box C



Patent family members are listed in annex

*** Special categories of cited documents**

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

- *X* document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

Z document member of the same patent family

Date of the actual completion of the international search

28 November 1997

Date of mailing of the international search report

16.12.97

Name and mailing address of the ISA

European Patent Office P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31 70) 340 2040, Tx 31 651 epo nl,
Fax (+31 70) 340 3016

Authorized officer

Yeats, S

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/BE 97/00092

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	D.L. BECK ET AL.: "Disruption of virus movement confers broad-spectrum resistance against systemic infection by plant viruses with a triple gene block" PROC. NATL. ACAD. SCI. USA, vol. 91, 1994, pages 10310-10314, XP002025561 see abstract.	1
A	--- D. GILMER ET AL.: "Efficient cell-to-cell movement of beet necrotic yellow vein virus requires 3' proximal genes located on RNA 2" VIROLOGY, vol. 189, 1992, pages 40-47, XP002025562 cited in the application see abstract and Figure 1. -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/BE 97/00092

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9113159 A	05-09-91	FR 2658987 A	06-09-91
		FR 2658988 A	06-09-91
		AT 129745 T	15-11-95
		DE 69114275 D	07-12-95
		DE 69114275 T	13-06-96
		EP 0517833 A	16-12-92
		ES 2079647 T	16-01-96
